

Use of a Canine Gastrointestinal Olfactory Stimulant in a Shelter Setting

Abigail Anne Williams¹, Ian Cunningham¹, Terry E Brady¹, Sarah K Abood², Rachel Tinker-Kulberg¹, Kristen Dellinger^{1,3}, Melinda KM Goddard¹, Lee Robertson¹ and Anthony L Dellinger^{1,3*}

Abstract

A recent investigation linking olfactory triggers to a distinct reflex has highlighted the relationship between “sniffing” the ground and canine defecation. The work presaged the question as to whether dogs awaiting adoption would exhibit the same apparent scent mediation of final peristalsis in a kennel environment. In fact, this study has shown that olfactory gastrointestinal neurobiological pathways remained intact with rescue dogs. Further, use of an olfactory stimulant resulted in more rapid and predictable excretion behavior consistent with prototypical household routines upon adoption. In turn, the canine olfactory stimulant improved operational efficiencies in a shelter management protocol with faster, more reliable “dog walking.” That is, with less time spent literally waiting for dogs to relieve themselves in often stressful, unfamiliar and densely populated shelter settings, more time would be available for interaction and exercise so as to leverage limited resources. Ultimately, this strategy may prove helpful to millions of shelter dogs, improve adoption rates and retention, and reduce relinquishments associated with home soiling.

Introduction

Dogs in the shelter system

According to the American Society for the Prevention of Cruelty to Animals (ASPCA), approximately 3.3 dogs enter US shelters every year, frequently overwhelming the infrastructure, resources and the staff working to care for them [1]. Of these, an estimated 1.6 million are adopted annually, and the cost of their supplies (i.e., food, bowls, collars, leashes, toys, etc.) to shelters often exceeds available budgets [1]. Notably, this excludes veterinary care, which can range from hundreds of thousands to millions of dollars annually depending on the size of a facility; whereas, many shelters spay and neuter animals prior to adoption, which can help prevent unwanted litters and encourage adoption [2-4]. These overarching factors culminate with euthanasia for more than one-third (38%) of total US admissions and almost half (48%) of relinquished dogs to shelters; while nearly one in five (18.8%) of rehomed dogs are returned and half of these are subsequently euthanized [5]. The interest in rescue adoptions has gradually increased, but with only 20-23% of Americans actually doing so, individual dog breeders still represent the greatest portion of new animal acquisitions at 34% [6,7]. While some shelters allow same-day adoptions, many organizations follow more rigorous procedures to ensure suitable placements; and practices can vary in relation to a given shelter's resources. With half of American households owning some 78 million dogs [8], the entire annual population of shelter dogs would nonetheless represent about 4% of the total, suggesting more potential growth for first or additional dog ownership in the US given higher rates of successful adoptions.

Effect of shelter setting on canine elimination behavior

Multiple studies have concluded that canines defecate and urinate away from the spaces in which they sleep, drink and eat whenever possible [9-11]. Despite millennia of domestication, canines likewise typically defecate most readily when outside in a safe, acceptable space when given the time to do so. The sensitivity and specificity of canine olfaction have also been well-documented [12]. As such, chemosensory information exchanges from external stimuli and the canine brain appear to be linked to the physiology of the autonomic and somatic nervous systems, suggesting association with both final peristalsis (voluntarily releasing the bowel) and prompting of defecation [13]. The retention and subsequent release of gut contents following universal “sniffing” in advance of defecation yields compelling evidence of such a connection in the canine olfactory bulb; this underscores the importance of exposure to olfactory stimulants whether located successfully from a “sniff search” on open grounds or when provided to supplement the dog's natural environment, whenever shelter staff time limits the ability for systematic exercise and “outdoor” access, it would follow that reducing the time associated with routine outdoor defecation could ease the burden on shelter staff sufficiently to help establish more predictable canine elimination habits, consistent with adoption expectations. Staff working in tandem could also use the time while the dogs are walked outside for cleaning and disinfecting crates or cages to minimize the animals' exposure to cleaning agents and reduce the potential for transmission of pathogens and parasites throughout the shelter.

Canine olfaction and gastrointestinal neurobiology

Whether living as pets or awaiting adoption at a shelter, many dogs have been observed to “sniff” for extended periods and seemingly ignore competing scents throughout their “search” prior to relieving themselves. This behavior has been described as the Recto-Anal Inhibitory Reflex (RAIR) or rectal distention in the absence of specific olfactory stimulation. RAIR allows the animal to delay defecation by moving the stool inward slightly and reducing the urge to defecate [14]. Add in the vagaries of indoor disinfectants and efforts to keep outdoor kennel runs clean and the “sniff search” for specific scents could also lead to RAIR due to the scarcity of such molecules. While some pet owners may more readily indulge such delays, this aspect of canine behavior can thwart shelter efforts to maintain healthy exercise and excretion routines with limited staffing. In turn, upon olfactory stimulation, dogs can activate voluntary muscles to initiate movement of the stool forward and outside of the body. Such behavior links the canine's “sniff search” to act as a precursor to relief and further suggests that particular organic scents induce these reflexes to facilitate elimination [15]. In turn, use of a stimulant solution formulated with such naturally occurring molecules could likewise leverage available staff time if more dogs could be “walked” and/or sequentially given “outdoor” access with the same resources due to more predictable and timely defecation by each dog.

Methods

Shelter description and operation

To test the potential canine behavioral and operational management impact of an olfactory stimulant to expedite defecation in a shelter setting, canine participants were selected in collaboration with a regional facility [SPCA of the Triad, Brown Summit, NC (SPCAT)]. The SPCAT maintains approximately 50 dogs of various ages and breeds at any given time and usually operates with one staff member and a volunteer at opening. These individuals are responsible for moving larger dogs to outdoor enclosures, cleaning cages, replacing bedding material and toys, taking the smaller dogs outside for walks to eliminate, and preparing food and water for each animal. The dogs are crated and secured inside the facility from 6:00PM to 6:00-7:00 AM for 12 to 13 hours, 7 days a week.

Selection of canine participants

Consent was obtained by shelter directors; the staff's elected the canine participants; and field-testing was supervised by the resident kennel team. In order to be included in the study, the dogs needed to be healthy, capable of being walked with a collar and leash, in the care of the SPCAT, and estimated to be at least 6 months of age. Recruitment was ongoing, as new animals arrived at the shelter throughout the 8-week trial. The total time that each animal had been held in the shelter varied across dogs, with some new arrivals and others having been kenneled for more than two years. Dogs included in the study ranged from approximately 1 to 11 years of age, and both males and females were evaluated. In all, 25 shelter animals were selected for evaluation; however, many were adopted prior to full assessment, leaving 14 of them accessible for the entire trial. A summary of breeds is shown in Table 1.

Baseline and treatment assessment

Throughout the study, the animals were crated indoors without access to food or water each night. The minimum assessment criteria required at least four timed baseline walks, followed by at least seven timed treatment walks. The baseline average untreated defecation times (TD) were determined without application of the olfactory stimulant. For the treated walks, a single drop of stimulant solution was placed on the front paw, a timer was started, and the dog was walked around the outdoor property (~1.2 acres). The timer was stopped at the onset of elimination. If dogs eliminated in the crate overnight, their times were not recorded, except for those of one that did so repeatedly: A 3-year old male Beagle mix nonetheless produced stool during the timed walks, which were included in the trial data. Typically, larger dogs that were located outside during operating hours were walked first, followed by the smaller dogs that were kept indoors in crates throughout the day. Each assessment began in the morning at approximately 6:30 AM and typically concluded by 8:30 AM. The canines' elimination behavior was evaluated by beginning a timer after each dog was leashed, brought outside and walked around the designated outdoor space. The timer was stopped at the onset of each defecation and the time was recorded. In addition to elimination time, stool quality, animal behavior, and other environmental observations were recorded for each test. Each dog was then additionally walked for 5 to 10 minutes for exercise before returning to their indoor or outdoor kennel for the day: Larger dogs (n=4) were maintained outside in individual cages (~6 × 10') and smaller dogs (n=10) remained indoors in crates (~2 × 3').

Breed	Age (Yrs)	Gender	Time in the Shelter	Spayed/Neutered
Aussie mix	2	Female	<3 months	Yes
Australian cattle mix	4	Female	>18 months	Yes
Carolina dog mix	3	Female	>18 months	Yes
Chihuahua	2	Female	<3 months	Yes
Chihuahua	1	Female	<1 month	Yes
Pitbull terrier mix	3	Female	>18 months	Yes
Shih Tzu	11	Female	<3 months	Yes
Basset hound Labrador mix	2	Male	>6 months	Yes
Beagle mix	3	Male	>18 months	Yes
Flat-coated retriever mix	1	Male	<3 months	Yes
Hound mix	7.5	Male	<3 months	Yes
Labrador mix	9	Male	>18 months	Yes
Shepherd Pitbull mix	2	Male	>6 months	Yes
Spitz mix	3	Male	<3 months	Yes

Table 1: Breed, age, and gender of evaluated shelter dogs (n=14).




Figure 1: Field Trials with 1 to 2 Year-old Shelter Canines: Aussie Mix, Basset Hound Labrador Mix, Chihuahuas (2), Flat-coated Retriever Mix, and Shepherd Pitbull Mix. A) The average untreated defecation time (T₀; red bars) for each dog was established individually with at least 4 observations over a 1-week period. Gray “striped” bars represent the average treated T₀ using an olfactory scent stimulant with >7 evaluations. B) Indicates details, average untreated/defecation times, total treatment evaluations, and percent change from untreated vs. treated T₀ of each dog. Significant differences (p<0.05; Student's t-test) between treated and untreated T₀ indicated by (*).




Figure 2: Field Trials with 3 to 4 Year-old Shelter Canines: Australian Cattle Mix, Beagle, Carolina Dog Mix, Pitbull Terrier Mix, and Spitz Mix. A) The average untreated defecation time (T₀; red bars) for each dog was established individually with at least 4 observations over a 1-week period. Gray “striped” bars represent the average treated T₀ using an olfactory scent stimulant with >7 evaluations. B) Indicates details, average untreated/defecation times, total treatment evaluations, and percent change from untreated vs. treated T₀ of each dog. Significant differences (p<0.05; Student's t-test) between treated and untreated T₀ indicated by (*).







Figure 3: Field Trials with >4 Year-old Shelter Canines: Hound Mix and Labrador Mix. A) The average untreated defecation time (T₀; red bars) for each dog was established individually with at least 4 observations over a 1-week period. Gray “striped” bars represent the average treated T₀ using an olfactory scent stimulant with >7 evaluations. B) Indicates details, average untreated/defecation times, total treatment evaluations, and percent change from untreated vs. treated T₀ of each dog. Significant differences (p<0.05; Student's t-test) between treated and untreated T₀ indicated by (*).



Discussion

Discussion In an earlier study using an olfactory stimulant with domestic dogs in residential settings, approximately 75% of them demonstrated more rapid and consistent defecation; whereas, each of the participating dogs evaluated in the shelter setting (100%) responded with decreased defecation times in this study [13-15]. These findings suggest that the organic amines required to mediate canine gastrointestinal neurobiology may be literally harder to find in the relatively sterile and continually disinfected environment required to maintain a clean and safe shelter. Whereas, more of the previously evaluated pets had presumably acclimated to more fruitful “sniff searching” in their backyards or familiar grounds, where such naturally occurring organic compounds may have been more prevalent. In fact, the defecation response to botanical scents in both cohorts has further refuted the common misperception that dogs seek prior fecal deposits to stimulate defecation; this is distinctly different behavior than repetitive urination associated with competitive “marking” of a given territorial perimeter, as in one common example [26].

Epidemiologically, once impounded in a shelter, dogs face nearly a 50% probability of euthanasia; if adopted and returned to a shelter for house soiling, the odds may double. Simply put, shelters represent the front line of homeless dog and cat survival, where staff and volunteers might be compared to “first responders” that perform heroics, much as firefighters and police officers. If routine use of an olfactory stimulant can help sheltered dogs establish predictable and more rapid defecation behaviors, it would follow that administering precisely the same olfactory stimulant to which they responded in the shelter would likewise help orient them to unfamiliar household circumstances upon adoption. In turn, continued home use would enable new owners to help prompt appropriate defecation when let outside to a yard or taken for a walk with far greater probability of the same response in keeping with busy human schedules; thereby, reduce the risk of home soiling and the attendant consequences of one or more failed adoptions for any given animal.

Conclusion

Whether acquired from shelters or breeders, such human routines have also continued to shape canine daily life with more owners dependent on increasingly accessible, organized dog walking services; as apps and providers merge the virtual and real resources to address the exercise and elimination needs of the “home alone” pet. The business model often requires a dog walker to manage multiple canines simultaneously. Thus, the “sniff search” can be complicated by numerous distractions and represents the next exploration of canine olfactory mediation of gastrointestinal neurobiology, or the use of such a stimulant, in complex circumstances.

References

- American Society for the Prevention of Cruelty to Animals (2015) Pet Statistics.
- Cashion, Bettie W. What Does it Cost to Run a Shelter?
- Brother Wolf Animal Rescue (2019) Adoption Process.
- Animal Humane Society (2019) How Adoption Works.
- Patronek GJ, Glickman LT, Meyer MR (1995) Population dynamics and the risk of euthanasia for dogs in an animal shelter. *Anthrozoos* 8: 31-43.
- Mohan-Gibbons H, Weiss E, Garrison L, Allison M (2014) Evaluation of a Novel dog adoption program in two US communities. *PLoS One* 9: e91959.
- Scarlett JM, Salman MD, New Jr JG, Kass PH (1999) Reasons for relinquishment of companion animals in U.S. animal shelters: Selected health and personal issues. *J Appl Anim Welfare Science* 2: 41-57.
- Grueter LM, Barber RT, Wynne CD (2015) A canine identity crisis: Genetic breed heritage testing of shelter dogs. *PLoS one* 13: e0202633.
- Tynes VV (2007) Canine Houstraining Challenges. *Veterinary Medicine*.
- Wagner D, Newbury S, Kass P, Harley K (2014) Elimination behavior of shelter dogs housed in double compartment kennels. *PLoS one* 9: e96254.
- Newbury S, Binm MK, Bushby PA, Cox CB, Damage JD (2010) Facility design and environment. In *Guidelines for Standards of Care in Animal Shelters*.
- Brady TE, Abood SK, Kulberg RT, Dellinger K, Goddard MKM, et al. (2019) Olfactory mediation of canine gastrointestinal neurobiology. *J Anim Health Behav Sci* 3: 116-148.
- Andrews PL (1992) Physiology of nausea and vomiting. *Br J Anesth* 69: 2S-19S.
- Chen JH, Sallam HS, Lin L, Chen JD (2010) Colorectal and rectocolic reflexes in canines: involvement of tone, compliance and anal sphincter relaxation. *Am J Physiol Regul Integr Comp Physiol* 299: 953-959.
- Hirabayashi T, Matsufuji H, Yokoyama J, Hagane K, Hoshino K, et al. (2003) Colorectal motility induction by sacral nerve electrostimulation in a canine model. *Dis Colon Rectum* 46: 809-817.
- Hart BL (1974) Environmental and hormonal influences on urine marking behavior in the adult male dog. *Behavioral biology*, 11: 167-176.

Author Affiliations

- Kepley BioSystems Incorporated, 2901 East Gate City Blvd, Suite 2400, Greensboro, NC, USA, 27401
- Ontario Veterinary College, University of Guelph, 50 Stone Rd E., Guelph, ON, Canada, N1G2W1
- Joint School of Nanoscience and Nanoengineering, 2907 East Gate City Blvd, Greensboro, NC, USA, 27401

Williams AA, Cunningham I, Brady TE, Abood SK, Tinker-Kulberg R, et al. (2019) Use of a Canine Gastrointestinal Olfactory Stimulant in a Shelter Setting. *J Anim Health Behav Sci* 3: 117. [LINK to Article](#)